arXiv:0804.4352v1 [nucl-ex] 28 Apr 2008

Latest results from the PHOBOS experiment

Barbara Wosiek for the PHOBOS Collaboration ‡

Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland

E-mail: `barbara.wosiek@ifj.edu.pl`

B.Alver[4], B.B.Back[1], M.D.Baker[2], M.Ballintijn[4], D.S.Barton[2], R.R.Betts[6], A.A.Bickley[7], R.Bindel[7], W.Busza[4], A.Carroll[2], Z.Chai[2], V.Chetluru[6], M.P.Decowski[4], E.García[6], T.Gburek[3], N.George[2], K.Gulbrandsen[4], C.Halliwell[6], J.Hamblen[8], I.Harnarine[6], M.Hauer[2], C.Henderson[4], D.J.Hofman[6], R.S.Hollis[6], R.Hołyński[3], B.Holzman[2], A.Iordanova[6], E.Johnson[8], J.L.Kane[4], N.Khan[8], P.Kulinich[4], C.M.Kuo[5], W.Li[4], W.T.Lin[5], C.Loizides[4], S.Manly[8], A.C.Mignerey[7], R.Nouicer[2], A.Olszewski[3], R.Pak[2], C.Reed[4], E.Richardson[7], C.Roland[4], G.Roland[4], J.Sagerer[6], H.Seals[2], I.Sedykh[2], C.E.Smith[6], M.A.Stankiewicz[2], P.Steinberg[2], G.S.F.Stephans[4], A.Sukhanov[2], A.Szostak[2], M.B.Tonjes[7], A.Trzupek[3], C.Vale[4], G.J.van Nieuwenhuizen[4], S.S.Vaurynovich[4], R.Verdier[4], G.I.Veres[4], P.Walters[8], E.Wenger[4], D.Willhelm[7], F.L.H.Wolfs[8], B.Wosiek[3], K.Woźniak[3], S.Wyngaardt[2], B.Wysłouch[4]

[1] Argonne National Laboratory, Argonne, IL 60439-4843, USA

[2] Brookhaven National Laboratory, Upton, NY 11973-5000, USA
[3] Institute of Nuclear Physics PAN, Kraków, Poland
[4] Massachusetts Institute of Technology, Cambridge, MA 02139-4307, USA
[5] National Central University, Chung-Li, Taiwan
[6] University of Illinois at Chicago, Chicago, IL 60607-7059, USA
[7] University of Maryland, College Park, MD 20742, USA
[8] University of Rochester, Rochester, NY 14627, USA

Abstract. Over the past years PHOBOS has continued to analyze the large datasets obtained from the first five runs of the Relativistic Heavy Ion Collider (RHIC) at Brookhaven National Laboratory. The two main analysis streams have been pursued. The first one aims to obtain a broad and systematic survey of global properties of particle production in heavy ion collisions. The second class includes the study of fluctuations and correlations in particle production. Both type of studies have been performed for a variety of the collision systems, covering a wide range in collision energy and centrality. The uniquely large angular coverage of the PHOBOS detector and its ability to measure charged particles down to very low transverse momentum is exploited. The latest physics results from PHOBOS, as presented at Quark Matter 2008 Conference, are contained in this report.

---

‡ For the full list of PHOBOS authors and acknowledgments, see appendix "Collaborations"

PACS numbers: 25.75.-q

1. Introduction

PHOBOS [1] is a heavy ion experiment at the BNL RHIC collider, known for its capability of measuring charged particles over a broad angular acceptance, by far the largest of all RHIC experiments. With the PHOBOS multiplicity array charged particles are measured in pseudorapidity range $\mid \eta \mid < 5.4$ and over 2π in the azimuth. In addition, the unique design of the two-arm spectrometer allows for extending the particle momentum measurements to the lowest limit reachable at RHIC. With this detector we have collected data on p+p, d+Au, Cu+Cu and Au+Au collisions during the RHIC 2000-2005 runs, covering a wide range of collision energy and centrality. This comprehensive data set allowed for a systematic studies of the overall features of particle production mechanisms in nuclear and elementary collisions, which constituted the main part of the baseline PHOBOS physics program. While this part is now nearly completed, our current effort is mainly focused on the study of fluctuations and correlations in particle production, the study of which can provide deeper insight into different stages of the system evolution.

In this paper we briefly summarize the results on global properties of charged particle production, including antiparticle to particle ratios and particle yields at very low transverse momentum (p_T). The emphasis is put on our recent results obtained from fluctuation and correlation studies. The dynamical fluctuations of the elliptic flow, corrected for the non-flow effects are presented for Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV. The same high-statistics data set is used to investigate structures in the near- and away-side correlations with respect to high-p_T trigger particle over a broad range in $\Delta\eta$. Finally, the results from a systematic study of the two-particle angular correlations in p+p, Cu+Cu and Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV are shown.

2. System-size dependence of particle production

Recently PHOBOS has completed a systematic study of the bulk properties of the produced particles, like total multiplicity, $dN_{ch}/d\eta$ and dN_{ch}/dp_T distributions, particle composition and collective flow effects, in Cu+Cu and Au+Au collisions as a function of the collision energy and centrality. The results, in comparison to d+Au and elementary p+p collisions, show that particle production in heavy ion collisions can be described in terms of simple scaling rules. Number of participant (N_{part}) scaling is observed for the total particle multiplicity [2], net-proton yields [3] and also, as will be shown later, the low-p_T particle yields. Energy and centrality dependencies of mid-rapidity yields factorize over an extended range of transverse momenta [4, 5]. Particle yields and directed (v_1) and elliptic flow (v_2) signals measured over a wide range of high to almost central pseudorapidities show energy-independence when viewed in the rest frame of one of the colliding nuclei [2]. The antiparticle to particle ratios show at most weak dependence on the system size [6], as is illustrated by the compilation of PHOBOS data presented in Figure 1.



Figure 1: Antiparticle to particle ratios for pions, kaons and protons as a function of collision centrality for different collision systems, measured at mid-rapidity region of $0.2 < \eta < 1.4$.

It is also observed that N_{part} unifies the measurements of the nuclear modification factors in Cu+Cu and Au+Au collisions [7]. A consistent and unified description of the elliptic flow measured in Cu+Cu and Au+Au collisions can be obtained after scaling v_2 with the participant eccentricity, ϵ_{part} [8].

The above observations show that the collision geometry has a major impact on the dynamical evolution of the system. They also provide a tool for extrapolating RHIC data to the LHC energy regime, which is of particular importance since to date no theory or model can consistently explain the scaling rules controlling the particle production in heavy ion collisions.

3. Particle production at very low transverse momenta

The PHOBOS spectrometer design provides a unique capability of studying particle production at the lowest transverse momenta accessible at RHIC. The new results on low-p_T yields for pions, kaons and protons measured in a high-statistics sample Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV have been shown at this Conference [9]. Figure 2 shows the invariant low-p_T yields measured for 6% of the most central Au+Au collisions in comparison to the PHENIX data at higher p_T [10]. The low-p_T yields agree with the extrapolations of Blast-Wave and Bose-Einstein fits to the PHENIX results. A flattening of the shape of the p_T spectra is observed, stronger for heavier particles, which is consistent with a rapid transverse expansion of the system. Invariant yields, integrated over the low-p_T range ($0.031 - 0.053$ GeV/c for pions, $0.105 - 0.128$ GeV/c for kaons and $0.143 - 0.206$ GeV/c for protons and antiprotons) and normalized per

N_{part}, are shown as a function of N_{part} for Au+Au collisions at 200 and 62.4 GeV [3] in Figure 3. One can see that, within the errors, the invariant low-p_T yields scale with N_{part}. For more details see Ref.[9].



Figure 2: Identified particle p_T spectra near mid-rapidity in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV.



Figure 3: The integrated low-p_T yields for Au+Au collisions at 200 GeV (solid symbols) and 62.4 GeV (open symbols) as a function of centrality.

4. Elliptic flow fluctuations

The measured magnitude of the elliptic flow signal, v_2, should reflect the initial spatial eccentricity of the overlap region of the colliding nuclei. As PHOBOS has shown [8], the participant eccentricity, ϵ_{part}, which takes into account the fluctuations in the participant nucleon positions in the overlap region, is the relevant eccentricity which drives the observed azimuthal anisotropy. We have used the Monte Carlo Glauber (MCG) approach to study the robustness of the participant eccentricity and also to calculate the higher order cumulants of ϵ_{part} quantifying the event-by-event fluctuations of the initial source eccentricity [11]. It was shown that the participant eccentricity is a robust quantity, insensitive to the choice of MCG parameters and model assumptions and that it strongly fluctuates with $\sigma(\epsilon_{part})/\langle\epsilon_{part}\rangle$ varying from about 35% for peripheral up to 50% for most central Au+Au collisions.

Comparison of the v_2 measurements to hydrodynamic model predictions [12] indicate that the matter created in heavy ion collisions at RHIC has properties resembling those of a perfect fluid [13]. In hydrodynamic models, the fluctuations in the shape of the initial collision geometry should be reflected in the fluctuations of v_2. PHOBOS has measured large (of the order of $40-50\%$) relative fluctuations of the elliptic flow, $\sigma(v_2)/\langle v_2\rangle$ for Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV [14]. The main difficulty in this measurement is to disentangle dynamical fluctuations of v_2 and non-flow correlations due, for example, to resonance decays or jet production. PHOBOS has developed a new method to extract the non-flow component from the data. The method relies on the determination of the second Fourier coefficient of two-particle angular correlations measured over a broad range of $\Delta\eta$, thanks to the large acceptance of the PHOBOS detector. This coefficient, $v_2^2(\eta_1, \eta_2)$, contains contributions from genuine flow correlations, $v_2(\eta_1) \times v_2(\eta_2)$, and non-flow correlations, $\delta(\eta_1, \eta_2)$. At $\Delta\eta > 2$, we assume that the non-flow component is small. In fact a small non-flow effect at $\Delta\eta > 2$ was estimated from HIJING simulations and accounted for. Thus, performing the fit to the second Fourier coefficient of two-particle angular correlations allows us to determine the flow component, $v_2(\eta_1) \times v_2(\eta_2)$, and then estimate the non-flow contribution by subtracting flow component from $v_2^2(\eta_1, \eta_2)$. The contribution from non-flow correlations to $\sigma(v_2)$ is $\sqrt{\langle\delta\rangle/2}$ [15]. Figure 4 shows the measured total relative v_2 fluctuations and the non-flow contribution to these fluctuations. The non-flow contribution extracted from the data is large, of the order of $25 - 30\%$. The relative flow fluctuations, after subtracting the non-flow contribution, are shown in Figure 5 and compared to the relative fluctuations of the initial eccentricity calculated from MCG [11] and Color Glass Condensate (CGC) [16] models. It can be seen that the magnitude of the relative flow fluctuations agree, within the errors, with both MCG and CGC calculations of the fluctuations in initial source eccentricity, leaving essentially no room for other, later-stage contributions. For more details see Ref. [17].



Figure 4: Total measured relative flow fluctuations (circles) and the contribution from non-flow correlations (squares) for Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV. Bands show 90 % C.L. systematic errors.



Figure 5: Relative flow fluctuations, corrected for the non-flow effects, compared to the relative fluctuations of the initial eccentricity calculated from MC Glauber and CGC models.

5. High-p_T triggered two-particle correlations

Measurements of correlations with respect to the high-p_T trigger particle allow us to study the medium response to energetic partons produced in early hard scattering processes and then propagated through the dense medium created in heavy ion collisions. Here we report the first measurements of the $\Delta\eta$ and $\Delta\phi$ correlations between the triggered charged particle with $p_T > 2.5$ GeV/c and all associated hadrons recorded in the PHOBOS Octagon detector [1]. The analysis utilizes the extensive coverage in pseudorapidity ($|\ \eta\ |< 3$) and azimuthal angle ($\Delta\phi = 2\pi$) of the Octagon allowing for the study of both short- and long-range correlations in $\Delta\eta$. A particular emphasis is put on testing the presence of long-range correlations at near- and away-sides over the uniquely broad acceptance in $\Delta\eta$ ($-4 < \Delta\eta < 2$). The correlated yields per trigger particle, with the elliptic flow effects subtracted, are calculated for Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV and compared to the correlated particle production in p+p events simulated with PYTHIA. The two dimensional correlated yield, $\frac{1}{N_{trig}}\frac{d^2N_{ch}}{d\Delta\eta d\Delta\phi}$, in central Au+Au collisions exhibits a much broader away-side peak in $\Delta\phi$ as compared to p+p correlations. The near-side yield in central Au+Au collisions shows a clear jet-like peak near $\Delta\phi \approx 0$ as well as pronounced correlations extending over a broad $\Delta\eta$ range (the so called ridge structure). In Figure 6 the correlated near-side yield (integrated over $|\ \Delta\phi\ |< 1$) is plotted as a function of $\Delta\eta$ for 10% of the most central Au+Au collisions. A prominent ridge effect is clearly visible up to $\Delta\eta = -4$ in contrast to p+p PYTHIA events, where the structure is absent. The momentum kick model [18] is able to quantitatively describe this structure. Centrality dependence of the near-side yield, averaged over $-4 < \Delta\eta < -2$, is shown in Figure 7. One can see that the ridge effect weakens with decreasing collision centrality and almost completely disappears for N_{part} less than about 80. For more details of this analysis see Ref. [19].



Figure 6: The correlated near-side yield for 0-10% central Au+Au collisions compared to p+p PYTHIA simulations (dashed line) and momentum kick model predictions (solid line).



Figure 7: The near-side yield averaged over $-4 < \Delta\eta < -2$ as a function of N_{part}.

6. Two-particle angular correlations

The study of the correlations between the final-state particles should provide additional information on the underlying mechanism of particle production at freeze-out. PHOBOS has performed systematic studies of two-particle angular correlations in $(\Delta\eta, \Delta\phi)$ in elementary p+p collisions [20] as well as Cu+Cu collisions [21]. Recently these studies have been also carried out for Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV. The broad coverage of the PHOBOS Octagon detector is fully utilized in these studies. The multiplicity-independent two-particle correlation function, $R(\Delta\eta, \Delta\phi)$, is calculated using the method detailed in [20]. We have concentrated on studying the short-range correlations by projecting the two-dimensional correlation function into a one-dimensional correlation function $R(\Delta\eta)$. $R(\Delta\eta)$ exhibits clear short-range correlations which can be described using a simple cluster model. The one-dimensional correlation function is fitted to a parameterization derived in an independent cluster model [22] in order to extract the effective cluster size, K_{eff}. Figure 8 shows the effective cluster size as a function of N_{part} for $\sqrt{s_{NN}} = 200$ GeV Cu+Cu and Au+Au collisions. For both collision systems the cluster size decreases with increasing centrality. Furthermore, for the same N_{part}, clusters in Au+Au collisions are larger than in the Cu+Cu system. Interestingly, this dependence on the size of the colliding nuclei disappears after plotting the same data as a function of fractional cross-section as shown in Figure 9. More differential studies of the effective size of near- and away-side clusters have also been performed. It is observed that away-side clusters are smaller and decrease more rapidly with increasing centrality than near-side clusters. This observation can be qualitatively explained by assuming that near-side clusters are preferentially produced close to the surface of the emission zone, while away-side clusters seem to propagate through the medium. For more details on the two-particle correlation study see [23].



Figure 8: Effective cluster size as a function of N_{part} for Cu+Cu and Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV. Dashed line denotes the value of K_{eff} measured in p+p collisions.



Figure 9: The same as in Figure 8 but plotted against the fractional cross-section.

7. Summary and outlook

PHOBOS is continuing to provide interesting and unique data on particle production in different collision systems in the RHIC energy range. The uniqueness of the PHOBOS data is owed to the distinct features of the PHOBOS detector: its extensive acceptance for charged particle measurements and capability of measuring particles at very low p_T. Comprehensive results on global features of the particle production are only briefly summarized in this report. The measurements of low-p_T yields, for the first time performed as a function of centrality, are presented for Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV. No anomalous enhancement of pion production is observed at very low p_T. The spectral shapes for heavier particles show flattening effects, consistent with a strong transverse expansion of the system.

In this report we have presented a wealth of results on the fluctuations and correlation measurements from PHOBOS. New results on event-by-event elliptic flow fluctuations, corrected for the non-flow effects extracted from data, are shown for Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV. Although non-flow correlations contribute significantly to the measured flow fluctuations, the corrected relative flow fluctuations are large, with a magnitude in agreement with calculations of fluctuations in the participant eccentricity. These results indicate that system thermalizes very rapidly and the initial-state event-by-event source fluctuations are efficiently converted into final-state momentum fluctuations.

The PHOBOS measurements of correlations between the trigger high-p_T particle and associated particles with no p_T cut imposed, allow for studying the ridge structure in the near-side correlations over the longitudinal extent larger than accessible in other RHIC experiments. The results for Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV show that the ridge structure of the near-side correlations persists up to the limit of our acceptance, i.e. $\Delta\eta = 4$. With decreasing collision centrality the structure becomes less pronounced, eventually disappearing at N_{part} of about 80. These results provide valuable constraints on models aimed to describe jet propagation through a dense medium.

The systematic studies of two-particle angular correlations for p+p, Cu+Cu and Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV show that at freeze-out particles tend to be produced in clusters with a non-trivial centrality dependence observed in nucleus-nucleus collisions. Furthermore, an unexpected system size dependence is seen. These observations provide a challenge to models describing particle production in heavy ion collisions.

The PHOBOS will continue the analysis of the existing comprehensive dataset. The studies of the low-p_T particle production as well as of elliptic flow fluctuations and triggered correlations will be extended to other systems and energies. Although PHOBOS has accomplished its initial goal, the future studies should still provide results improving our understanding of the physics of heavy ion collisions.

This work was partially supported by U.S. DOE grants DE-AC02-98CH10886, DE-

FG02-93ER40802, DE-FG02-94ER40818, DE-FG02-94ER40865, DE-FG02-99ER41099, and DE-AC02-06CH11357, by U.S. NSF grants 9603486, 0072204, and 0245011, by Polish MNiSW grant N N202 282234 (2008-2010), by NSC of Taiwan Contract NSC 89-2112-M-008-024, and by Hungarian OTKA grant (F 049823).

References

[1] Back B B *et al.* (PHOBOS) 2003 *Nucl. Instrum. Meth.* **A 499** 603.
[2] Alver B *et al.* (PHOBOS) arXiv:0709.4008 [nucl-ex].
[3] Back B B *et al.* (PHOBOS) 2007 *Phys. Rev.* **C 75** 024910.
[4] Roland G *et al.* (PHOBOS) 2006 *Nucl. Phys.* **A 774** 113.
[5] Hofman D J *et al.* (PHOBOS) 2007 *J. Phys.* **G 34** s217.
[6] Alver B *et al.* (PHOBOS) arXiv:0802.1695 [nucl-ex], submitted to *Phys. Rev.C.*
[7] Alver B *et al.* (PHOBOS) 2006 *Phys. Rev. Lett.* **96** 212301.
[8] Alver B *et al.* (PHOBOS) 2007 *Phys. Rev. Lett.* **98** 242302.
[9] Gburek T *et al.* (PHOBOS), these proceedings.
[10] Adler S S *et al.* (PHENIX) 2004 *Phys. Rev.* **C 69** 034909.
[11] Alver B *et al.* (PHOBOS) 2008 *Phys. Rev.* **C 77** 014906.
[12] Teaney D 2003 *Phys. Rev.* **C 68** 034913.
[13] Kovtun P, Son D T and Starinets A O 2005 *Phys. Rev. Lett.* **94** 111601.
[14] Alver B *et al.* (PHOBOS) 2007 arXiv:0702036 [nucl-ex].
[15] Voloshin S A, Poskanzer A M, Tang A, Wang G 2008 *Phys. Lett.* **B 659** 537.
[16] Drescher H J and Nara Y 2007 *Phys. Rev.* **C 76** 041903.
[17] Alver B *et al.* (PHOBOS), these proceedings.
[18] Wong C Y, private communications, see also these proceedings.
[19] Wenger E *et al.* (PHOBOS), these proceedings.
[20] Alver B *et al.* (PHOBOS) 2007 *Phys. Rev.* **C 75** 054913.
[21] Li W *et al.* (PHOBOS) 2007 *J. Phys.* **G 34** s1005.
[22] Morel A and Plaut G 1974 *Nucl. Phys.* **B 78** 541.
[23] Li W *et al.* (PHOBOS), these proceedings.